

## K-Ar AGES OF IGNEOUS AND METAMORPHIC ROCKS FROM MCMURDO, SØR RONDANE AND DUMONT D'URVILLE, ANTARCTICA

Naoko UENO<sup>1</sup>, Atsumi OGATA<sup>2</sup> and Minoru FUNAKI<sup>3</sup>

<sup>1</sup>*Toyo University, 11-10, Oka 2-chome, Asaka 351*

<sup>2</sup>*Institute for Study of the Earth's Interior,  
Okayama University, Misasa, Tottori 682-01*

<sup>3</sup>*National Institute of Polar Research, 9-10, Kaga 1-chome, Itabashi-ku, Tokyo 173*

**Abstract:** K-Ar radiometric ages were determined for igneous and metamorphic rocks from the Antarctic Margin. Twenty-five volcanics of 11 sites from McMurdo Sound resulted in between 0.08-Ma and 4.15-Ma. Two syenites from the Sør Rondane Mountains were about 400 Ma and three granitic rocks from Dumont d'Urville were around 1000 Ma.

### 1. Introduction

Many types of fresh lithologies with ages from Archean to Quaternary have been found in Antarctica, and have attracted many earth scientists, in particular, paleomagnetists who are interested in continental drift and/or paleointensity of earth magnetism. The paleomagnetic studies should be done with rock samples of which the ages have been determined with radiometric dating methods. In this paper, the authors will present K-Ar ages for representative igneous and metamorphic rocks from McMurdo Sound, Sør Rondane Mountains and Dumont d'Urville where FUNAKI (1984) and others have carried out paleomagnetic studies.

### 2. Geology and Samples

#### 2.1. McMurdo Sound

Volcanism occurred sporadically in this region in the Plio-Pleistocene (MANKINEN and COX, 1988). The representative volcanic ejecta were collected from Hut Point Peninsula and Cape Royds on Ross Island and from Taylor Valley (Fig. 1). The sampling sites on the Hut Point Peninsula are illustrated in Fig. 2. Petrographic descriptions for all the samples are listed in the Appendix. MANKINEN and COX (1988) reported paleomagnetic data combined with previously published geologic and geochronologic data, and determined the general sequences of volcanic rocks of the McMurdo volcanic province. Previous reports have shown K-Ar ages of Black Knob to be  $0.43 \pm 0.07$  Ma (ARMSTRONG, 1978), Cape Royds to be  $0.68 \pm 0.14$  Ma (TREVES, 1967), Half Moon Crater to be  $1.0 \pm 0.15$  Ma (ARMSTRONG, 1978), Castle Rock to be  $1.1 \pm 0.4$  Ma (KYLE and TREVES, 1974), and Observation Hill to be  $1.18 \pm 0.03$  Ma (FORBES *et al.*, 1974), respectively. ARMSTRONG (1978)




Fig. 1. A topographic map of the McMurdo Sound region and the sampling sites (modified after FUNAKI, 1984).




Fig. 2. Sampling sites of Hut Point Peninsula (modified after FUNAKI, 1984).

reported K-Ar ages (1.5 Ma to 4.6 Ma) of rocks from Taylor Valley, and gave the ages  $2.93 \pm 0.10$  Ma and  $2.87 \pm 0.15$  Ma for the basaltic rocks. We also collected basaltic rocks from the same location as ARMSTRONG (1978) did.

## 2.2. Sør Rondane Mountains

The Sør Rondane Mountains in East Antarctica consist of medium- to high-grade metamorphic rocks, and various plutonic and minor dike rocks (SHIRAISHI and KAGAMI, 1992). GREW (1982) compiled radiometric age data in this region; the gneisses have U-Pb zircon ages of 2700 Ma, 950–970 Ma and 520–600 Ma, and the plutonic rocks have only zircon age of 520–600 Ma. TAKIGAMI and FUNAKI (1991) reported  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  and K-Ar ages of K-feldspar and biotite from igneous and metamorphic rocks and suggested igneous activity of 440–500 Ma and metamorphism of 350–400 Ma. They also reported a K-Ar biotite age of  $662 \pm 23$  Ma from gneiss. SHIRAISHI and KAGAMI (1992) obtained Sm-Nd and Rb-Sr whole rock isochron ages around 1000 Ma for granulite facies metamorphic rocks,  $624 \pm 18$  Ma Sm-Nd internal mineral isochron age of hornblende, biotite and plagioclase, and  $556 \pm 26$  and  $483 \pm 12$  Ma Rb-Sr mineral ages of hornblende, biotite and plagioclase. Sampling sites in the Sør Rondane Mountains are illustrated in Fig. 3. Petrographic descriptions of the samples are shown in the Appendix.

## 2.3. Dumont d'Urville

The Dumont d'Urville region is situated around the Wilkes Subglacial Basin, which trends north, parallel to the transatlantic mountains, and has exposure of high-grade metamorphic rocks of Proterozoic. GREW (1982) reported Rb-Sr biotite ages from pegmatite and granite of 1530 and 1543 Ma, respectively. Samples were collected from Ile Cuvier near Ile du Lion and from Ile des Petrels (Fig. 4). Petrographic descriptions of the samples are shown in the Appendix.




Fig. 3. Sampling sites in the Sør Rondane Mountains (modified after TAKIGAMI and FUNAKI, 1991).




Fig. 4. Sampling sites in Dumont d'Urville (modified after Institut Geographique National en 1954).

### 3. Experiments

K content was analysed by flame photometry at Toyo University using a 1000 ppm Cs buffer. The detailed analytical method was described in UENO (1991). Experimental error of the K content was assumed to be 5% based on multiple analyses of the standard samples (JB-2, JB-3 and JG-1a by Geological Survey of Japan described in ANDO *et al.*, 1989).

Ar analysis was performed by a noble gas mass spectrometer (Modified VG5400) installed in the Institute for Study of the Earth's Interior, Okayama University. About 0.1–0.5 g of sieved samples between 250  $\mu\text{m}$  and 180  $\mu\text{m}$  were used for analyses of McMurdo Sound rocks, which were expected to be within 0.5 and 3.0 Ma. About 50 mg of samples were analysed for other regions where the rock ages were expected to be older than 500 Ma. Samples were wrapped with Al foil 8  $\mu\text{m}$  thick and put into a sample holder. The samples from McMurdo Sound were set at once with a working standard sample YZ1-6 ( $0.236 \pm 0.004$  Ma with K content of 1.41 wt%, *cf.* NAGAO *et al.*, 1991), separately from other old samples which were set up with a working standard sample MMhb-1 ( $517.5 \pm 4.5$  Ma with K content of 1.58 wt%, *cf.* SAMSON and ALEXANDER, 1987). Analyses were carried out in a computer controlled peak jumping mode. Details of the analytical technique and the age calculation will be reported elsewhere (NAGANO *et al.*, in preparation).

### 4. Results and Discussion

The results of K-Ar analyses are shown in Tables 1, 2 and 3. The errors of K-Ar ages

Table 1. Summary of K-Ar ages in McMurdo Sound.

	Rock type	K (Wt %)	Weight (g)	$^{36}\text{Ar}$ ( $10^{-10} \text{ cm}^3 \text{ STP/g}$ )	$^{40}\text{Ar}/^{36}\text{Ar}$	$^{40}\text{Ar}_{\text{rad}}$ ( $10^{-8} \text{ cm}^3 \text{ STP/g}$ )	Age (Ma)	$^{40}\text{Ar}_{\text{fair}}/^{40}\text{Ar}_{\text{total}}$ (%)
Cape Royds	Kenite	3.35	0.5498	$3.956 \pm 0.072$	$322.96 \pm 0.64$	$1.067 \pm 0.030$	$0.082 \pm 0.005$	91.7
	Kenite	3.52	0.4830	$6.911 \pm 0.101$	$311.81 \pm 0.57$	$1.093 \pm 0.041$	$0.080 \pm 0.005$	94.9
	Kenite	3.23	0.5582	$5.456 \pm 0.083$	$314.47 \pm 0.58$	$1.008 \pm 0.033$	$0.080 \pm 0.005$	94.1
	Kenite	3.15	0.5626	$5.422 \pm 0.082$	$313.66 \pm 0.54$	$0.957 \pm 0.031$	$0.078 \pm 0.005$	94.4
Taylor Valley	Ankaramite	1.28	0.1544	$22.721 \pm 0.267$	$358.90 \pm 0.74$	$14.29 \pm 0.22$	$2.87 \pm 0.15$	82.5
	Ankaramite	1.19	0.1516	$16.745 \pm 0.226$	$410.06 \pm 1.25$	$19.18 \pm 0.25$	$4.15 \pm 0.21$	72.2
Hut point Peninsula								
Crater Hill	Ankaramite	1.11	0.2238	$22.813 \pm 0.257$	$303.19 \pm 0.51$	$1.641 \pm 0.114$	$0.39 \pm 0.04$	97.6
	Ankaramite	1.25	0.2561	$4.645 \pm 0.089$	$343.72 \pm 1.06$	$2.216 \pm 0.060$	$0.46 \pm 0.03$	86.1
Twin Crater	Ankaramite	1.24	0.2576	$5.723 \pm 0.097$	$336.40 \pm 0.83$	$2.312 \pm 0.057$	$0.48 \pm 0.03$	88.0
	Ankaramite	1.20	0.2016	$5.627 \pm 0.114$	$333.78 \pm 1.00$	$2.126 \pm 0.066$	$0.46 \pm 0.03$	88.7
Between Cape	Ankaramite	1.35	0.2122	$3.829 \pm 0.096$	$355.55 \pm 1.67$	$2.280 \pm 0.078$	$0.44 \pm 0.03$	83.2
Arm. & Scott B.	Ankaramite	1.32	0.2363	$16.376 \pm 0.193$	$313.34 \pm 0.53$	$2.840 \pm 0.089$	$0.55 \pm 0.03$	94.5
North of	Olivine basalt	1.63	0.2435	$6.956 \pm 0.106$	$359.79 \pm 1.06$	$4.437 \pm 0.090$	$0.70 \pm 0.04$	82.3
Hut Point	Olivine basalt	1.55	0.2198	$7.025 \pm 0.113$	$352.08 \pm 0.97$	$3.940 \pm 0.085$	$0.65 \pm 0.04$	84.1
	Olivine basalt	1.52	0.2904	$20.141 \pm 0.236$	$353.31 \pm 0.76$	$11.54 \pm 0.18$	$1.95 \pm 0.10$	83.8
Hut Point	Olivine basalt	1.46	0.2200	$16.241 \pm 0.194$	$320.20 \pm 0.59$	$3.930 \pm 0.099$	$0.69 \pm 0.04$	92.4
Castle Rock	Olivine basalt	1.55	0.2217	$15.529 \pm 0.187$	$324.74 \pm 0.62$	$4.463 \pm 0.103$	$0.74 \pm 0.04$	92.2
	Olivine basalt	1.57	0.2178	$29.984 \pm 0.331$	$310.38 \pm 0.64$	$4.312 \pm 0.188$	$0.71 \pm 0.05$	95.4
Black Knob	Olivine basalt	1.30	0.3942	$0.916 \pm 0.088$	$461.88 \pm 12.34$	$1.514 \pm 0.020$	$0.30 \pm 0.02$	64.1
	Olivine basalt	1.29	0.3614	$5.278 \pm 0.111$	$371.08 \pm 1.41$	$3.971 \pm 0.062$	$0.79 \pm 0.04$	79.8
Half Moon	Hornblende basalt	3.38	0.2340	$16.450 \pm 0.193$	$352.94 \pm 0.90$	$9.367 \pm 0.164$	$0.71 \pm 0.04$	83.9
Crater	Hornblende basalt	1.91	0.2101	$10.144 \pm 0.175$	$365.34 \pm 1.20$	$7.033 \pm 0.155$	$0.95 \pm 0.05$	81.0
Obs. Hill	Trachyte	3.73	0.2011	$18.245 \pm 0.218$	$399.67 \pm 0.92$	$18.95 \pm 0.25$	$1.31 \pm 0.07$	74.1
	Trachyte	3.57	0.2552	$17.482 \pm 0.198$	$496.40 \pm 2.13$	$17.48 \pm 0.20$	$1.26 \pm 0.06$	59.6
	Trachyte	3.28	0.3100	$12.064 \pm 0.168$	$428.57 \pm 1.47$	$16.05 \pm 0.20$	$1.26 \pm 0.06$	69.1

Analytical errors for K concentrations were estimated to be 5%.  $\pm$  indicates one standard deviation. \*Whole rock samples removed magnetite by hand magnet were used for analyses. \*\*Whole rock samples were used for analyses.

are represented as one standard deviation.

#### 4.1. *McMurdo Sound*

Samples have been collected from three separate volcanic regions ; Cape Royds, Hut Point Peninsula and Taylor Valley (Fig. 1). They are all very fresh except for some samples which show oxidation texture (Nos. 78012203, 78012205, 77121508, 78122310), and have partly altered olivine (No. 77121607). Olivine has no potassium, suggesting that the partial alteration has no significant effect on the K-Ar age of the whole rock sample. Oxidation occurs in general in clinker part of flowing lava. The samples with oxidation texture also present no critical problem in K-Ar analysis of rocks older than 1 Ma, though this type of sample sometimes has unusual argon isotope systematics (OKADA *et al.*, 1991).

The four kenytes from Cape Royds are about 0.08 Ma, indicating that the volcanic activity in this area continued at least from 0.68 Ma, as reported by TREVES (1967) up to the Latest Pleistocene.

The two ankaramites from Taylor Valley are  $2.87 \pm 0.15$  and  $4.15 \pm 0.21$  Ma in age, indicating volcanic activity with time span longer than 1.2 million years in the region. The younger age ( $2.87 \pm 0.15$ ) is similar to the ages reported by ARMSTRONG (1978), who dated the rocks below Marr Glacier where we collected this sample, while the older one is closer to the age ( $4.64 \pm 0.12$  and  $4.5 \pm 0.7$  Ma) of the dikes east of Solas Glacier in Taylor Valley by ARMSTRONG (1978).

Table 2. Summary of K-Ar ages in the Sør Rondane Mountains.

	Rock type	K (wt %)	Weight (g)	$^{36}\text{Ar}$ ( $10^{-10} \text{ cm}^3$ STP/g)	$^{40}\text{Ar}/^{36}\text{Ar}$	$^{40}\text{Ar}_{\text{rad}}$ ( $10^{-4} \text{ cm}^3$ STP/g)	Age (Ma)	$^{40}\text{Ar}_{\text{air}}/^{40}\text{Ar}_{\text{total}}$ (%)
B749	Syenite	8.18	0.0513	$27.504 \pm 0.657$	$49826 \pm 1749$	$1.432 \pm 0.033$	$*402 \pm 20$	0.6
B799	Syenite	7.70	0.0490	$24.466 \pm 0.429$	$49629 \pm 1567$	$1.310 \pm 0.026$	$*392 \pm 19$	0.6
B859	Basalt	1.84	0.0603	$18.888 \pm 0.328$	$32458 \pm 0709$	$0.636 \pm 0.008$	$**724 \pm 30$	0.9

Analytical errors for K concentrations were estimated to be 5%.

$\pm$  indicates one standard deviation.

Whole rock samples removed magnetite by hand magnet were used for analyses.

\*K-feldspar is partly altered to clay minerals.

\*\*Thermally metamorphosed sample. The calculated age is without geological significance.

Table 3. Summary of K-Ar ages at Dumont d'Urville.

	Rock type	K (wt %)	Weight (g)	$^{36}\text{Ar}$ ( $10^{-10} \text{ cm}^3$ STP/g)	$^{40}\text{Ar}/^{36}\text{Ar}$	$^{40}\text{Ar}_{\text{rad}}$ ( $10^{-4} \text{ cm}^3$ STP/g)	Age (Ga)	$^{40}\text{Ar}_{\text{air}}/^{40}\text{Ar}_{\text{total}}$ (%)
E-1	Syenite	7.40	0.0533	$47.705 \pm 0.732$	$69431 \pm 3278$	$3.354 \pm 0.209$	$0.900 \pm 0.057^*$	0.4
E-225	Quartz monzonite	5.61	0.0531	$31.253 \pm 0.816$	$93757 \pm 6400$	$3.073 \pm 0.175$	$1.04 \pm 0.06^*$	0.3
E-323	Granite	3.26	0.0528	$31.921 \pm 0.842$	$68518 \pm 3288$	$2.314 \pm 0.095$	$1.26 \pm 0.06^*$	0.4

Analytical errors for K concentrations were estimated to be 5%.

$\pm$  indicates one standard deviation.

Whole rock samples were used for analyses.

\*K-feldspar is partly altered to clay mineral.

In the Hut Point Peninsula region, four types of volcanic rocks, ankaramite, olivine basalt, hornblende basalt and trachyte, were collected from nine volcanic domains. Ankaramites from Crater Hill, Twin Crater, and between Cape Armitage and Scott Base have ages varying from 0.39 to 0.55 Ma, belonging to the youngest volcanics in the region. Trachytes from Observation Hill are about 1.3 Ma in age, the oldest in the region. Basalts from other domains have an intermediate age range from about 0.7 to 0.95 Ma, except for two extremely old ( $1.95 \pm 0.10$  Ma) and young ( $0.30 \pm 0.02$  Ma) rocks from Hut Point and Black Knob.

#### 4.2. Sør Rondane Mountains

The two syenites (Nos. B749, B799) from Lunckeryggen show whole rock K-Ar ages of 402 and 392 Ma, respectively. These ages are slightly younger than the total fusion biotite and K-feldspar age (413–498 Ma) of the  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  method by TAKIGAMI and FUNAKI (1991) who collected syenites (Nos. B740, B788) from the same domain.

The basalt (No. B859) from Brattinipene was thermally metamorphosed in microscopic observation. The calculated age of 724 Ma of this sample might be overestimated, and is without geological significance.

#### 4.3. Dumont d'Urville

The syenite (No. E1), granite (No. E323) and quartz monzonite (No. E225) are 0.900, 1.26 and 1.04 Ga in age respectively. The ages are significantly younger than the Rb-Sr biotite age of about 1.5 Ga found by GREW (1982). This might also be also due to rejuvenation by alteration because these samples have some secondary minerals.

### 5. Summary

We obtained the K-Ar ages of the igneous and metamorphic rocks from the Antarctic Margin as follows.

- (1) In McMurdo Sound, new K-Ar data have been added, ranging from 0.08 to 4.15 Ma.
- (2) For the Sør Rondane Mountains, about 400 Ma has been obtained from partly altered samples.
- (3) For Dumont d'Urville, 0.90–1.26 Ga is obtained from partly altered samples.

### Acknowledgments

Prof. K. NAGAO of the Institute for Study of the Earth's Interior, Okayama University supported the Ar analyses. We owe microscopic observation and sample descriptions to Dr. T. TIBA of the National Science Museum. The unknown referees, Prof. I. KANEOKA of Tokyo University and Prof. Y. TAKIGAMI of Kanto Gakuen University gave comments to improve the manuscript.

### References

AANDO, A., MITA, N. and MATSUMOTO, A. (1989): 1987 compilation of  $\text{K}_2\text{O}$  concentrations in seventeen

- GSJ rock reference samples, "Igneous rock series". Bull. Geol. Surv. Jpn., **40**, 19-45.
- ARMSTRONG, R.L. (1978): K-Ar dating: Late Cenozoic McMurdo volcanic group and Dry Valley glacial history, Victoria Land, Antarctica. N. Z. J. Geol. Geophys., **21**, 685-698.
- FORBES, R.B., TURNER, D.E. and CARDEN, J.R. (1974): Age of trachyte from Ross Island, Antarctica. Geology, **2**, 297-298.
- FUNAKI, M. (1984): Paleomagnetic investigation of McMurdo Sound region, Southern Victoria Land, Antarctica. Mem. Natl Inst. Polar Res., Ser. C (Earth Sci.), **16**, 1-81.
- GREW, E.S. (1982): The Antarctic Margin. The Oceans Basins and Margins, Vol. 6, The Indian Ocean, ed. by A.E.M. NAIRN and F.G. STEHLI. New York, Plenum, 697-755.
- KYLE, P.R. and TRVES, S.B. (1974): Geology of Hut Point Peninsula, Ross Island. Antarct. J. U. S., **9**, 232-234.
- MANKINEN, E.A. and COX, A. (1988): Paleomagnetic investigation of some volcanic rocks from the McMurdo volcanic province, Antarctica. J. Geophys. Res., **93**, 11599-11612.
- NAGAO, K., OGATA, A., MIURA, Y., MATSUDA, J. and AKIMOTO, S. (1991): Highly reproducible 13 and 17 ka K-Ar ages of two volcanic rocks. Geochem. J., **25**, 447-451.
- OKADA, T., ITAYA, T., SAWA, I. and HONG, S.H. (1991): K-Ar determination of a lava stalagmite in Manjang gul, Jeju island, Korea. J. Southeast Asian Earth Sci., **6**, 127-130.
- SAMSON, S. and ALEXANDER, E.C., Jr. (1987): Calibration of the inter laboratory  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  dating standard, MMhb-1. Chem. Geol. (Isot. Geosci. Sec.), **66**, 27-34.
- SHIRAIISHI, K. and KAGAMI, H. (1992): Sm-Nd and Rb-Sr ages of metamorphic rocks from the Sør Rondane Mountains, East Antarctica. Recent Progress in Antarctic Earth Science, ed. by Y. YOSHIDA *et al.* Tokyo, Terra Sci. Publ., 29-35.
- TAKIGAMI, Y. and FUNAKI, M. (1991):  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  and K-Ar ages for igneous and metamorphic rocks from the Sør Rondane Mountains, East Antarctica. Proc. NIPR Symp. Antarct. Geosci., **5**, 122-135.
- TREVES, S.B. (1967): Volcanic rocks from the Ross Island, Marguerite Bay and Mt. Weaver areas, Antarctica. JARE Sci. Rep., Spec. Issue, **1**, 136-149.
- UENO, N. (1991): Ganseki hyôjun shiryô JB-2, JB-3 oyobi JG-1a-chû no K no teiryô (Potassium concentrations in GSJ rock reference samples-JB-2, JB-3 and JG-1a). Tôyô Daigaku Kiyô Kyôyôkatei-hen (Shizen Kagaku) (J. Toyo Univ. Gen. Educ. (Nat. Sci.)), **35**, 1-10.

(Received March 1, 1994; Revised manuscript received June 17, 1994)

*Appendix. Rock types and constituent minerals (after Tokiko TIBA).*

Abbreviations: ph: phenocryst gr: groundmass ap: apatite bi: biotite  
 cpx: clinopyroxene gl: glass hm: hematite ho: hornblende  
 il: ilmenite mt: magnetite ol: olivine op: opaque mineral  
 pl: plagioclase qz: quartz

---

McMurdo Sound

- 79010903: Kenyte, fresh  
 ph: anorthoclase with dusty inclusions and inclusions of cpx, op.  
 gr: pl, anorthoclase, cpx, op, bi.
- 79010940: Kenyte, fresh  
 ph: anorthoclase with dusty inclusions and inclusions of sodic  
 cpx, ol, cpx, op, ap. gr: pl, anorthoclase, op, bi, cpx.
- 79010944: Kenyte, fresh  
 ph: anorthoclase with inclusions of cpx and ap and dusty  
 inclusions, sodic augite, op, ol. gr: pl (sodic), cpx, op.
-



*Appendix (continued).*


---

79011009 :	Kenyte, fresh ph : anorthoclase with inclusions of op and cpx and dusty inclusions. gr : pl (sodic), cpx, op.
77121607 :	Ankaramite ph : ol-partly altered to iddingsite, cpx. gr : pl, cpx, op, ol.
77112704 :	Ankaramite, fresh ph : ol, cpx. gr : pl, cpx, op, brown gl.
77112707 :	Ankaramite, fresh ph : ol, cpx. gr : pl, cpx, op, brown gl.
77112711 :	Ankaramite, fresh ph : ol, cpx, pl, op. gr : pl, cpx, op, brown gl.
78012104 :	Ankaramite, fresh ph : ol, cpx. gr : pl, cpx, op, brown gl.
78012105 :	Ankaramite, fresh ph : ol, cpx. gr : pl, cpx, op, ol ?, brown gl.
78011805 :	Olivine basalt, fresh ph : cpx, ol. gr : pl, cpx, op (mt, il), brown gl, ol ?, accretional lapilli.
78011806 :	Olivine basalt, fresh ph : ol, cpx. gr : pl, cpx, op (mt, il), brown gl, ol ?, mesostases-cpx, il, hm, mt, brown gl.
77121602 :	Olivine basalt, fresh ph : ol, cpx. gr : pl, ol, cpx, op (mt, il), brown gl.
77121603 :	Olivine basalt, fresh ph : ol, cpx. gr : pl, cpx, ol, op, brown gl.
77121910 :	Olivine basalt, porous, fresh microph : ol, op. gr : pl, cpx, ol ?, op, brown gl, hm.
77121913 :	Olivine basalt, porous, fresh microph : ol, cpx. gr : pl, cpx, ol ?, op (mt, il), bi, gl.
78012214 :	Olivine basalt, fresh ph : ol, cpx. gr : pl, cpx, op (mt, il), brown gl.
78012215 :	Basalt, fresh ph : cpx, ol, pl. gr : pl, cpx, ol, op.
78012203 :	Hornblende basalt (oxidized, pre-eruption) ph : brown ho (opacitized), opx ?, cpx. gr : pl, cpx, mt (partly oxidized to hm), hm, ap.
78012205 :	Hornblende basalt (oxidized, pre-eruption) ph : brown ho (opacitized), cpx. gr : pl, cpx, hm, opx, gl.
77121508 :	Trachyte (oxidized, pre-eruption) microph : ho (opacitized). gr : pl (sodic), ho, op, gl.
77121509 :	Trachyte (oxidized, pre-eruption) microph : ho (opacitized), anorthoclase. gr : pl, ho, op, gl.
78122310 :	Trachyte (oxidized, pre-eruption) microph : ho (opacitized), anorthoclase. gr : anorthoclase, ho, op, bi.
77122102 :	Ankaramite ph : ol-partly oxidized along cracks, cpx. gr : pl, op, hm, cpx.
79010408 :	Ankaramite, porous, fresh ph : ol, cpx. gr : pl, cpx, op, hm, ol (mantled with op grains).

---

*Appendix (continued).*


---

Sør Rondane Mountains	
B749:	Syenite major : K-feldspar with exsolved patches and bi inclusions in preferred orientation and partly altered to clay mineral, pl, green ho, bi, qz. minor : titanite, carbonite (calcite), ap, zircon, op.
B799 :	Syenite major : K-feldspar with exsolved patches and bi inclusions in preferred orientation and partly altered to clay mineral, pl, green ho, qz, bi. minor : ap, titanite, zircon, op.
B859 :	Basalt, thermally metamorphosed ph : cpx, opx mantled with cpx, pl. gr : pl, bi, cpx, smectite, op.

---

Dumont d'Urville	
E1 :	Syenite major : K-feldspar with dusty inclusion, exsolved patches, inclusions of hematite, partly altered to clay mineral. minor : ho, bi, op, zircon.
E225 :	Quartz monzonite major : K-feldspar with dusty inclusion and partly altered to clay mineral, pl, qz, bi. minor : op, zircon.
E323 :	Granite major : K-feldspar partly altered to clay mineral, qz, pl. minor : op, bi, ho, zircon, titanite, secondary actinolite, clay mineral.

---